

Unless you have studied chemistry at some point in your life, you probably would not know what alkenes are. However, if I were to tell you that alkenes are used to make plastic, you would probably have a better idea. Alkenes also have properties that are similar to alkanes, which makes them suitable fuels as well. Today, we shall discuss what alkenes are and some of the reactions involving alkenes.

What are Alkenes?

As mentioned in [Organic Chemistry - An Introduction](#), alkenes are hydrocarbons with a general formula of C_nH_{2n} . To be more specific, alkenes are saturated aliphatic hydrocarbons, which means that they only contain carbon and hydrogen atoms. This makes alkenes similar to alkanes. Covalent bonds join adjacent carbon atoms. We have also mentioned that alkenes have a suffix -ene.


Here are some of the simpler alkenes that we commonly encounter.

n	Name	Molecular formula	Structural formula	M_r	Melting point / °C	Boiling point / °C	Physical state (r.t.p)
2	Ethene	C_2H_4	$CH_2=CH_2$	28	-169	-104	Gas
3	Propene	C_3H_6	$CH_3CH=CH_2$	42	-185	-48	Gas
4	Butene	C_4H_8	$CH_3CH_2CH=CH_2$	56	-185	-6	Gas
5	Pentene	C_5H_{10}	$CH_3(CH_2)_2CH=CH_2$	70	-165	30	Liquid
6	Hexene	C_6H_{12}	$CH_3(CH_2)_3CH=CH_2$	84	-140	63	Liquid


“Methene” does not exist since it only has one carbon atom. As such, there is no double bonds between carbon atoms to form an alkene.

Alkenes with three or more carbon atoms exhibit isomerism. This includes constitutional isomerism due to the arrangement of carbon atoms, the position of $C=C$ bond, and cis-trans isomerism due to the restricted rotation of atoms about the $C=C$ bond.

For example, butene has four isomers - but-1-ene ($CH_3CH_2CH=CH_2$), 2-methylpropene ($CH_2=C(CH_3)_2$), cis-but-2-ene, and trans-but-2-ene.



cis-2-butene



trans-2-butene

What are Cycloalkenes?


Cycloalkenes are saturated alicyclic hydrocarbons - this means that they have a ring of carbon atoms. Unlike alkenes, cycloalkenes have a general formula of C_nH_{2n-2} . They are technically different from alkenes and contain the prefix *cyclo-* to distinguish them from the aliphatic alkenes. This makes cycloalkenes similar to cycloalkanes.

Here are some of the simpler cycloalkanes that we often see.

n	Name	Molecular formula	Structural formula	Melting point / °C	Boiling point / °C	Physical state (r.t.p)
---	------	-------------------	--------------------	--------------------	--------------------	------------------------

3 Cyclopropene C_3H_4

40 -127 -33 Gas




4 Cyclobutene C_4H_6




54 -90 13 Gas

5 Cyclopentene C_5H_8



68 -94 49 Liquid


6 Cyclohexene C_6H_{10}

82 7 81 Liquid

Properties of Alkenes

As mentioned earlier, alkenes have similar physical properties as alkanes.

Melting and boiling points

As the number of carbons in the hydrocarbon chain increases, more energy is needed to overcome the stronger instantaneous dipole-induced dipole (id-id) interactions between molecules. As such, the boiling point of the straight-chained alkane isomers increases. Therefore, the first three alkenes are gases at room temperature, whereas subsequent alkenes are liquids then solids. Likewise, this is similar to alkanes.

Branched-chained alkene isomers often have lower melting points than straight-chained alkene isomers. This is because they have less surface area of contact between molecules since they are more spherical. As such, there is a decrease in the strength of id-id interactions.

Cis alkene isomers are polar molecules as they have a larger net dipole moment than their

non-polar trans counterparts. As such, there is strong permanent dipole-permanent dipole (pd-pd) interactions between molecules. This gives rise to higher boiling points as more energy is needed to break the stronger pd-pd interactions. However, trans alkene isomers have higher melting points as they are more closely packed.

Other physical properties

Alkenes are colourless, non-polar molecules. They are insoluble in polar solvents but are soluble in non-polar solvents. This is due to the energy released between water molecules and alkene molecules being insufficient to overcome the [strong hydrogen bonds](#) between water molecules.

On top of that, alkenes are less dense than water. They also have stronger smells as compared to alkanes.

Preparing Alkenes

There are three ways to prepare alkenes – catalytic cracking of petroleum, dehydration of alcohols, and dehydrohalogenating halogenoalkanes. While catalytic cracking will produce a mixture of alkanes and alkenes, the other methods will only produce alkenes.


Catalytic Cracking of Petroleum

Catalytic cracking involves breaking down hydrocarbon chains into smaller alkanes and alkenes with a catalyst at a temperature of 600°C. The catalyst used is usually aluminium oxide or silicon (IV) oxide.

Cracking is used to produce petroleum fractions in demand, such as petrol and naphtha. On top of that, it can also produce hydrogen, which is needed in the Haber process to produce ammonia.


Dehydration of Alcohol

We can dehydrate alcohols in an elimination reaction under heat. However, excess concentrated sulfuric acid (H_2SO_4) or aluminium oxide (Al_2O_3) is needed.



Dehydrohalogenating Halogenoalkanes

Similar to the dehydration of alcohols, we can dehydrohalogenate halogenoalkanes in an elimination reaction under heat. In this case, either potassium hydroxide or sodium hydroxide in ethanol is used as a reactant. Two products will be formed - the major product (more substituted alkene) and the minor product (less substituted alkene). The major product has more alkyl groups bonded to the carbon atoms bound by the C=C bond.



Reactions

Electrophilic Addition


Pi electrons in the C=C bond are more exposed than sigma electrons, making them a good source of electrons. This causes the C=C bond to have a high electron density, making the alkene more susceptible to electrophile attacks. As such, alkenes are more reactive than alkanes.

That said, alkenes generally undergo electrophilic addition. This occurs when the weak pi bond in the C=C bond is broken to form two strong sigma bonds.

Reaction Reagents and Conditions

Mechanism


Symmetrical alkenes



Adding hydrogen halides


Gaseous hydrogen halides


Asymmetrical alkenes



Adding halogens in CCl_4 as solvent

Halogens in CCl_4 as solvent





Reduction

The reduction of alkenes form alkanes. For more information, read [our article on alkanes](#).

Oxidation

Reaction	Reagents and Conditions	Mechanism
Combustion	Oxygen	<p>In limited oxygen, alkenes may burn to give carbon monoxide and water. E.g. C₂H₄ (g) + 2O₂ (g) → 2CO (g) + 2H₂O (l)</p> <p>In excess oxygen, alkenes burn to give carbon dioxide and water. E.g. C₂H₄ (g) + 3O₂ (g) → 2CO₂ (g) + 2H₂O (l)</p> <p>This is similar to the combustion of alkanes. However, alkenes burn with a more smoky flame due to a higher concentration of carbon atoms.</p>
Diol formation	Cold KMnO ₄ in aqueous NaOH	The C=C bond undergoes mild oxidation when there is no heating. This results in the breaking of the pi bond and forming of the diol.
Carboxylic acid, ketones and CO ₂ formation	KMnO ₄ in aqueous H ₂ SO ₄ , heat	The C=C bond undergoes strong oxidation when there is heat. This results in the cleaving of the pi bond, and then the cleaving of the sigma bond.